

# THE FIRST $L^p$ -COHOMOLOGY OF SOME FINITELY GENERATED GROUPS AND $p$ -HARMONIC FUNCTIONS

MICHAEL J. PULS

**ABSTRACT.** Let  $G$  be a finitely generated infinite group and let  $p > 1$ . In this paper we make a connection between the first  $L^p$ -cohomology space of  $G$  and  $p$ -harmonic functions on  $G$ . We also describe the elements in the first  $L^p$ -cohomology space of groups with polynomial growth, and we give an inclusion result for nonamenable groups.

## 1. INTRODUCTION

In this paper  $G$  will always be a finitely generated infinite group and  $S$  will always denote a symmetric generating set for  $G$ . Let  $M$  be a right  $G$ -module and let  $C^n(G, M)$  be the set of functions from  $G^n = G \times \cdots \times G$  ( $n$  copies) to  $M$ . We now have a chain complex

$$0 \longrightarrow M \xrightarrow{\delta_0} C^1(G, M) \xrightarrow{\delta_1} C^2(G, M) \xrightarrow{\delta_2} \cdots \xrightarrow{\delta_{n-1}} C^n(G, M) \xrightarrow{\delta_n} \cdots$$

where for  $n \geq 0$

$$(\delta_n f)(g_1, \dots, g_{n+1}) = (f(g_2, \dots, g_{n+1})) \cdot g_1 + \sum_{k=1}^n (-1)^k f(g_1, \dots, g_k g_{k+1}, \dots, g_{n+1}) + (-1)^{n+1} f(g_1, \dots, g_n),$$

where  $g_k g_{k+1}$  occupies the  $k$ th position of  $f$ . Each  $\delta_n$  is a linear map and a calculation shows  $\delta_n \delta_{n-1} = 0$ . The  $n$ th group cohomology of  $G$  with coefficients in  $M$  is denoted by  $H^n(G, M)$  and is equal to  $\ker \delta_n / \text{im } \delta_{n-1}$ , where  $\ker \delta_n$  denotes the kernel of  $\delta_n$  and  $\text{im } \delta_{n-1}$  is the image of  $\delta_{n-1}$  in  $C^n(G, M)$ . If  $M$  is a Banach space then we can give  $C^n(G, M)$  the compact open topology. This means  $f_k \rightarrow f$  in  $C^n(G, M)$  if and only if  $f_k(g_1, \dots, g_n) \rightarrow f(g_1, \dots, g_n)$ . Note that  $\ker \delta_1$  is a Banach space. In general  $\text{im } \delta_{n-1}$  is not closed in  $\ker \delta_n$ . Let  $\overline{\text{im } \delta_{n-1}}$  denote the closure of  $\text{im } \delta_{n-1}$  in  $\ker \delta_n$ . The quotient space  $\overline{H}^n(G) = \ker \delta_n / \overline{\text{im } \delta_{n-1}}$  is called the  $n$ th reduced cohomology space of  $M$ .

Let  $\mathcal{F}(G)$  denote the set of real-valued functions on  $G$ . Let  $f \in \mathcal{F}(G)$  and let  $x \in G$ , the right translation of  $f$  by  $x$  is the function defined by  $f_x(g) = f(gx^{-1})$ . For a real number  $p \geq 1$ ,  $L^p(G)$  will consist of those  $f \in \mathcal{F}(G)$  for which  $\sum_{g \in G} |f(g)|^p < \infty$ , and  $C_0(G)$  will consist of those  $f \in \mathcal{F}(G)$  for which the set  $\{g \mid |f(g)| > \epsilon\}$  is finite for each  $\epsilon > 0$ . The sets  $\mathcal{F}(G)$ ,  $C_0(G)$  and  $L^p(G)$  are  $G$ -modules under right translations. In this paper we study  $H^1(G, L^p(G))$ , the first

---

*Date:* May 7, 2004.

*2000 Mathematics Subject Classification.* Primary: 43A15; Secondary: 20F65, 46N10.

*Key words and phrases.* nonamenable groups, groups with polynomial growth,  $L^p$ -cohomology,  $p$ -harmonic functions.

group cohomology of  $G$  with coefficients in  $L^p(G)$ . We also study  $\overline{H}_{(p)}^1(G)$ , the first reduced  $L^p$ -cohomology space.

This work was supported by a grant from the research office of Eastern Oregon University. The author would like to thank the university for their kind support.

## 2. PRELIMINARIES

Let  $\mathbb{R}G$  be the group ring of  $G$  over  $\mathbb{R}$ . For  $h \in \mathbb{R}G$  and  $f \in \mathcal{F}(G)$  we define a multiplication from  $\mathcal{F}(G) \times \mathbb{R}G$  into  $\mathcal{F}(G)$  by

$$(f * h)(g) = \sum_{x \in G} f(gx^{-1})h(x).$$

Observe that  $(f * (s - 1))(g) = f(gs^{-1}) - f(g)$  for each  $g \in G$  and each  $s \in S$ . For  $1 \leq p \in \mathbb{R}$ , let  $D^p(G) = \{f \in \mathcal{F}(G) \mid f * (s - 1) \in L^p(G) \text{ for each } s \in S\}$ . Recall that  $S$  is a symmetric set of generators for  $G$ . We define a norm on  $L^p(G)$  by  $\|f\|_p = (\sum_{g \in G} |f(g)|^p)^{\frac{1}{p}}$ , where  $f \in L^p(G)$ . Let  $h \in D^p(G)$  and let  $e$  be the identity element of  $G$ . We define a norm on  $D^p(G)$  by  $\|h\|_{D^p(G)} = (\sum_{s \in S} \|h * (s - 1)\|_p^p + |h(e)|^p)^{\frac{1}{p}}$ . Under this norm  $D^p(G)$  is a Banach space. Let  $f_1$  and  $f_2$  be elements of  $D^p(G)$ . We will write  $f_1 \simeq f_2$  if  $f_1 - f_2$  is a constant function. Clearly  $\simeq$  is an equivalence relation on  $D^p(G)$ . Identify the constant functions on  $G$  with  $\mathbb{R}$ . Now  $D^p(G)/\mathbb{R}$  is a Banach space under the norm induced from  $D^p(G)$ . That is, if  $[f]$  is an equivalence class from  $D^p(G)/\mathbb{R}$  then  $\|[f]\|_{D^p(G)/\mathbb{R}} = (\sum_{s \in S} \|f * (s - 1)\|_p^p)^{\frac{1}{p}}$ . We shall write  $\|f\|_{D(p)}$  for  $\|[f]\|_{D^p(G)/\mathbb{R}}$ . The norm for  $D^p(G)$  and  $D^p(G)/\mathbb{R}$  depends on the symmetric generating set  $S$ , but the underlying topology does not. If  $X \subset D^p(G)$ , then  $(\overline{X})_{D^p(G)}$  will denote the closure of  $X$  in  $D^p(G)$ . Similarly if  $Y \subset D^p(G)/\mathbb{R}$ , then  $(\overline{Y})_{D(p)}$  will denote the closure of  $Y$  in  $D^p(G)/\mathbb{R}$ . The cardinality of a set  $A$  will be denoted by  $|A|$ .

For the chain complex

$$0 \longrightarrow L^p(G) \xrightarrow{\delta_0} C^1(G, L^p(G)) \xrightarrow{\delta_1} C^2(G, L^p(G)) \xrightarrow{\delta_2} \dots$$

the map  $\delta_0$  is given by  $(\delta_0 f)(g) = f * (g - 1)$ . If  $f \in \mathcal{F}(G)$  and  $\delta_0 f \in C^1(G, L^p(G))$ , then  $\delta_1(\delta_0 f) = 0$  which implies  $D^p(G)/\mathbb{R} \subseteq \ker \delta_1$  since  $f * (g - 1) \in L^p(G)$  for all  $g \in G$ . We now show that the reverse inclusion is also true.

**Lemma 2.1.** *Let  $G$  be a finitely generated, infinite group. The kernel of  $\delta_1 : C^1(G, L^p(G)) \rightarrow C^2(G, L^p(G))$  is  $D^p(G)/\mathbb{R}$ .*

*Proof.* Consider the chain complex

$$0 \longrightarrow \mathcal{F}(G) \xrightarrow{\delta'_0} C^1(G, \mathcal{F}(G)) \xrightarrow{\delta'_1} C^2(G, \mathcal{F}(G)) \xrightarrow{\delta'_2} \dots$$

It was shown in [9, Lemma 4.2] that  $H^1(G, \mathcal{F}(G)) = 0$ . Thus for each  $f \in \ker \delta'_1$  there exists  $\bar{f}$  in  $\mathcal{F}(G)$  such that  $\delta'_0 \bar{f} = f$ .

Let  $f_1 \in C^1(G, L^p(G)) \subseteq C^1(G, \mathcal{F}(G))$  and suppose  $\delta_1 f_1 = 0$ . Then  $\delta'_1 f_1 = 0$  which implies there exists an  $\bar{f}_1 \in \mathcal{F}(G)$  such that  $\delta_0 \bar{f}_1 = f_1$ . The result now follows since  $\bar{f}_1 \in D^p(G)$ .  $\square$

The map  $\delta_0$  is an injection so we obtain the following:

- (a) The first cohomology group of  $G$  with coefficients in  $L^p(G)$ , denoted by  $H^1(G, L^p(G))$ , is isomorphic with  $D^p(G) / (L^p(G) \oplus \mathbb{R})$ .

- (b) The first reduced  $L^p$ -cohomology space of  $G$ , denoted by  $\overline{H}_{(p)}^1(G)$ , is isometric with  $D^p(G)/\overline{L^p(G)} \oplus \mathbb{R}$ , where the closure is taken in  $D^p(G)$ .

### 3. NONREDUCED $L^p$ -COHOMOLOGY AND $p$ -HARMONIC FUNCTIONS

Let  $f \in \mathcal{F}(G)$  and let  $g \in G$ . Suppose  $1 < p \in \mathbb{R}$  and define

$$\Delta_p f(g) := \sum_{s \in S} |f(gs^{-1}) - f(g)|^{p-2} (f(gs^{-1}) - f(g)).$$

In the case  $1 < p < 2$ , we make the convention that  $|f(gs^{-1}) - f(g)|^{p-2} (f(gs^{-1}) - f(g)) = 0$  if  $f(gs^{-1}) = f(g)$ . We shall say that  $f$  is  $p$ -harmonic if  $f \in D^p(G)$  and  $\Delta_p f(g) = 0$  for all  $g \in G$ . Recall that  $f$  is a harmonic function if  $\sum_{s \in S} (f(gs^{-1}) - f(g)) = 0$  for all  $g \in G$ . Let  $HD^p(G)$  be the set of  $p$ -harmonic functions on  $G$ . Observe that the constant functions are in  $HD^p(G)$ . If  $p = 2$ , then  $HD^2(G)$  is the linear space of harmonic functions on  $G$  with finite energy. In general,  $HD^p(G)$  is not a linear space if  $p \neq 2$ . A wealth of information about  $p$ -harmonic functions on graphs and manifolds can be found in [6, 7]. Many of the ideas in this section come from the paper [16]. In this section we will give a decomposition of  $D^p(G)$  that will allow us to make a connection between  $p$ -harmonic functions on  $G$  and  $\overline{H}_{(p)}^1(G)$ .

We begin by giving some preliminary results for both  $D^p(G)$  and  $HD^p(G)$ . Let  $f$  and  $h$  be elements in  $D^p(G)$  and let  $1 < p \in \mathbb{R}$ . Define

$$\begin{aligned} \langle \Delta_p h, f \rangle &:= \sum_{g \in G} \sum_{s \in S} |h(gs^{-1}) - h(g)|^{p-2} (h(gs^{-1}) - h(g)) (f(gs^{-1}) - f(g)) \\ &= \sum_{g \in G} \sum_{s \in S} |(h * (s - 1))(g)|^{p-2} ((h * (s - 1))(g)) ((f * (s - 1))(g)). \end{aligned}$$

The above sum exist since  $\sum_{g \in G} \sum_{s \in S} |h(gs^{-1}) - h(g)|^{p-2} (h(gs^{-1}) - h(g)) \|^q < \infty$ , where  $\frac{1}{p} + \frac{1}{q} = 1$ . The next lemma will be used to help show the uniqueness of the decomposition of  $D^p(G)$  that will be given in Theorem 3.5.

**Lemma 3.1.** *Let  $f_1$  and  $f_2$  be functions in  $D^p(G)$ . Then  $\langle \Delta_p f_1 - \Delta_p f_2, f_1 - f_2 \rangle = 0$  if and only if  $f_1 * (s - 1) = f_2 * (s - 1)$  for all  $s \in S$ .*

*Proof.* Let  $f_1, f_2 \in D^p(G)$  and assume there exists  $s \in S$  such that  $f_1 * (s - 1) \neq f_2 * (s - 1)$ . Define a function  $f : [0, 1] \rightarrow \mathbb{R}$  by  $f(t) = \sum_{g \in G} \sum_{s \in S} |f_1(gs^{-1}) - f_1(g) + t((f_2(gs^{-1}) - f_2(g)) - (f_1(gs^{-1}) - f_1(g)))|^p$ . Observe that  $f(0) = \|f_1\|_{D(p)}^p$  and  $f(1) = \|f_2\|_{D(p)}^p$ . A derivative calculation gives  $\frac{df}{dt}|_{t=0} = p \langle \Delta_p f_1, f_2 - f_1 \rangle = -p \langle \Delta_p f_1, f_1 - f_2 \rangle$ . It follows from Proposition 5.4 on page 24 of [3] that  $\|f_2\|_{D(p)} > \|f_1\|_{D(p)} - p \langle \Delta_p f_1, f_1 - f_2 \rangle$ . Similarly,  $\|f_1\|_{D(p)} > \|f_2\|_{D(p)} - p \langle \Delta_p f_2, f_2 - f_1 \rangle$ . Hence,  $p \langle \Delta_p f_1 - \Delta_p f_2, f_1 - f_2 \rangle > 0$  if there exists  $s \in S$  such that  $f_1 * (s - 1) \neq f_2 * (s - 1)$ .

Conversely, suppose  $f_1 * (s - 1) = f_2 * (s - 1)$  for all  $s \in S$ . Then  $\langle \Delta_p f_1 - \Delta_p f_2, f_1 - f_2 \rangle = 0$  since  $f_1 - f_2$  is a constant function on  $G$ .  $\square$

For  $g \in G$ , define  $\delta_g$  by  $\delta_g(x) = 0$  if  $x \neq g$  and  $\delta_g(g) = 1$ .

**Lemma 3.2.** *Let  $h \in \mathcal{F}(G)$ . Then  $h$  is a  $p$ -harmonic function if and only if  $\langle \Delta_p h, \delta_x \rangle = 0$  for all  $x \in G$ .*

*Proof.* Let  $x \in G$  and suppose  $h \in HD^p(G)$ , then

$$\langle \Delta_p h, \delta_x \rangle = -2 \sum_{s \in S} |h(xs^{-1}) - h(x)|^{p-2} (h(xs^{-1}) - h(x)) = 0.$$

Conversely, if  $\langle \Delta_p h, \delta_x \rangle = 0$  for all  $x \in G$ , then  $\sum_{s \in S} |h(xs^{-1}) - h(x)|^{p-2} (h(xs^{-1}) - h(x)) = 0$  for all  $x \in G$ .  $\square$

*Remark 3.3.* It follows immediately from the lemma that if  $h \in HD^p(G)$ , then  $\langle \Delta_p h, f \rangle = 0$  for all  $f \in \mathbb{R}G$ .

**Proposition 3.4.** *If  $h \in HD^p(G)$  and  $f \in (\overline{L^p(G)})_{D^p(G)}$ , then  $\langle \Delta_p h, f \rangle = 0$ .*

*Proof.* Let  $f \in (\overline{L^p(G)})_{D^p(G)}$  and let  $h \in HD^p(G)$ . There exists a sequence  $\{f_n\}$  in  $\mathbb{R}G$  such that  $\|f - f_n\|_{D^p(G)} \rightarrow 0$  as  $n \rightarrow \infty$ , since  $(\overline{\mathbb{R}G})_{D^p(G)} = (\overline{L^p(G)})_{D^p(G)}$ . Now

$$\begin{aligned} 0 &\leq |\langle \Delta_p h, f \rangle| \\ &= |\langle \Delta_p h, f - f_n \rangle| \\ &= \left| \sum_{g \in G} \sum_{s \in S} |(h * (s-1))(g)|^{p-2} ((h * (s-1))(g)) ((f - f_n) * (s-1))(g) \right| \\ &\leq \sum_{g \in G} \sum_{s \in S} |(h * (s-1))(g)|^{p-1} |((f - f_n) * (s-1))(g)| \\ &\leq \left( \sum_{g \in G} \sum_{s \in S} |((h * (s-1))(g))|^{p-1} \right)^q \|f_n - f\|_{D^p(G)} \rightarrow 0 \end{aligned}$$

as  $n \rightarrow \infty$ . The last inequality follows from Hölder's inequality  $\square$

We now give a decomposition of  $D^p(G)$  that will allow us to determine representatives for the nonzero classes in  $\overline{H}_{(p)}^1(G)$ .

**Theorem 3.5.** *Let  $1 < p \in \mathbb{R}$  and suppose  $\overline{L^p(G)}_{D^p(G)} \neq D^p(G)$ . If  $f \in D^p(G)$ , then we can write  $f = u + h$ , where  $u \in (\overline{\mathbb{R}G})_{D^p(G)}$  and  $h \in HD^p(G)$ . This decomposition is unique up to a constant function.*

*Proof.* Let  $f \in D^p(G)$  and let  $r$  equal the distance of  $f$  from  $(\overline{\mathbb{R}G})_{D^p(G)}$  in the  $D^p(G)$ -norm. Set  $B = \{v \in (\overline{\mathbb{R}G})_{D^p(G)} \mid \|f - v\|_{D^p(G)} \leq r + 1\}$ . Now  $B$  is a nonempty weakly compact set in the reflexive Banach space  $D^p(G)$ . The function  $F(v) = \|f - v\|_{D^p(G)}$  is a weakly lower semi-continuous function on  $B$ , so  $F(v)$  assumes a minimum value on  $B$ . This minimum must be  $r$ . Let  $u \in (\overline{\mathbb{R}G})_{D^p(G)}$  where  $\|f - u\|_{D^p(G)} = r$  and set  $h = f - u$ . We now proceed to show that  $h$  is  $p$ -harmonic on  $G$ . Let  $t \in \mathbb{R}$  and let  $w \in \mathbb{R}G$ . Now,  $\|h\|_{D^p(G)} = \|f - u\|_{D^p(G)} \leq \|f - (u - tw)\|_{D^p(G)}$  for all  $t \in \mathbb{R}$ . The minimum of  $\|f - (u - tw)\|_{D^p(G)}$  occurs when  $t = 0$ . Thus  $\frac{d}{dt} (\|h + tw\|_{D^p(G)})|_{t=0} = \sum_{g \in G} \sum_{s \in S} p |h(gs^{-1}) - h(g)|^{p-2} (h(gs^{-1}) - h(g)) (w(gs^{-1}) - w(g)) = 0$ . Using  $\delta_x$  for  $w$  in the above derivative calculation we obtain  $-2p \sum_{s \in S} |h(xs^{-1}) - h(x)|^{p-2} (h(xs^{-1}) - h(x)) = 0$ . Thus  $h$  is  $p$ -harmonic by Lemma 3.2.

We now show that this decomposition is unique up to a constant. Suppose  $f = u_1 + h_1 = u_2 + h_2$ , where  $u_1, u_2 \in (\overline{\mathbb{R}G})_{D^p(G)}$  and  $h_1, h_2 \in HD^p(G)$ . Now,  $\langle \Delta_p h_1 - \Delta_p h_2, h_1 - h_2 \rangle = \langle \Delta_p h_1 - \Delta_p h_2, u_2 - u_1 \rangle = 0$  by Proposition 3.4 since  $u_2 - u_1 \in (\overline{\mathbb{R}G})_{D^p(G)}$ . By Lemma 3.1,  $h_1 * (s-1) = h_2 * (s-1)$  for all  $s \in S$ .

Thus  $h_1 - h_2$  is a constant function, which implies that  $u_1 - u_2$  is also a constant function.  $\square$

We saw in section 2 that  $\overline{H}_{(p)}^1(G) = D^p(G)/\overline{L^p(G)} \oplus \mathbb{R}$ , so it follows from the theorem that each nonzero class in  $\overline{H}_{(p)}^1(G)$  can be represented by a nonconstant element in  $HD^p(G)$ . Thomas Schick has found an error in the proof of Theorem 5.3 from [10]. We were unable to fix the proof of that theorem using the techniques of [10]. The following corollary can be considered a partial fix of the theorem since many groups that have a central element of infinite order are also of polynomial growth.

**Corollary 3.6.** *If  $G$  is a finitely generated group with polynomial growth, then  $\overline{H}_{(p)}^1(G) = 0$  for  $1 < p \in \mathbb{R}$ .*

*Proof.* It was shown in [8, Corollary 1.10] that  $HD^p(G) = \mathbb{R}$  if  $G$  has polynomial growth. The result now follows.  $\square$

#### 4. NONAMENABLE GROUPS AND $L^p$ -COHOMOLOGY

In this section we will give some results concerning the first  $L^p$ -cohomology space of nonamenable groups.

Let  $A$  be a subset of a group  $G$  and define  $\partial A := \{x \in A \mid \text{there exists } s \in S \text{ with } xs \notin A\}$ . We shall say that a group  $G$  is *amenable* if it has an exhaustion  $G_1 \subset G_2 \subset \cdots, \cup_{k=1}^\infty G_k = G$  by finite subsets such that  $\lim_{k \rightarrow \infty} \frac{|\partial G_k|}{|G_k|} = 0$ . A group that is not amenable is said to be *nonamenable*. Our first result shows how amenability affects the way  $L^p(G)$  sits inside  $D^p(G)/\mathbb{R}$ .

**Theorem 4.1.** *Let  $G$  be a finitely generated infinite group and let  $1 \leq p \in \mathbb{R}$ . Then  $L^p(G)$  is closed in  $D^p(G)/\mathbb{R}$  if and only if  $G$  is nonamenable.*

*Proof.* Assume that  $G$  is nonamenable and let  $f \in (\overline{L^p(G)})_{D(p)} = (\overline{\mathbb{R}G})_{D(p)}$ . Now there exists a sequence  $\{f_n\}$  in  $\mathbb{R}G$  such that  $f_n \rightarrow f$  in the Banach space  $D^p(G)/\mathbb{R}$ . Thus  $\{f_n\}$  is a Cauchy sequence in  $D^p(G)/\mathbb{R}$ . It was shown in [4] that there exist a constant  $C_p$  such that  $\|u\|_p \leq C_p \|u\|_{D(p)}$  for all  $u \in \mathbb{R}G$  if and only if  $G$  is a finitely generated nonamenable group. Thus  $\{f_n\}$  is also a Cauchy sequence in  $L^p(G)$ . So there exists a  $\bar{f} \in L^p(G)$  such that  $\|\bar{f} - f_n\|_p \rightarrow 0$  as  $n \rightarrow \infty$ . Since  $L^p$ -convergence implies pointwise convergence we have that  $\|(\bar{f} - f_n) * (s - 1)\|_p \rightarrow 0$  as  $n \rightarrow \infty$  for each  $s \in S$ . Hence  $\|\bar{f} - f_n\|_{D(p)} \rightarrow 0$  as  $n \rightarrow \infty$ . Therefore,  $\bar{f} = f$ .

Conversely, suppose there exists an exhaustion  $G_1 \subseteq G_2 \subseteq \cdots, \cup_{k=1}^\infty G_k = G$  by finite subsets such that  $\lim_{k \rightarrow \infty} \frac{|\partial G_k|}{|G_k|} = 0$ . Let  $\chi_k$  denote the characteristic function on  $G_k$ . Now define a function  $f_k$  on  $G$  by  $f_k := \frac{\chi_k}{\sqrt[p]{|G_k|}}$ . Note that  $\|f_k\|_p = 1$  for all  $k$ . Computing the  $D^p(G)/\mathbb{R}$ -norm of  $f_k$  we obtain  $\|f_k\|_{D(p)}^p = \sum_{s \in S} \|f_k * (s - 1)\|_p^p \leq 2 \frac{|\partial G_k|}{|G_k|}$ . Thus,  $\lim_{k \rightarrow \infty} \|f_k\|_{D(p)} \rightarrow 0$  and  $\|f_k\|_p = 1$  for all  $k$ . Therefore, if  $G$  is amenable, then  $L^p(G)$  is not closed in  $D^p(G)/\mathbb{R}$ .  $\square$

The next result follows immediately from the theorem.

**Corollary 4.2.** *Let  $1 \leq p \in \mathbb{R}$ . If  $G$  is a finitely generated nonamenable group, then  $H^1(G, L^p(G)) = \overline{H}_{(p)}^1(G)$  and  $H^1(G, L^p(G))$  is a Banach space.*

We will now use the theorem to show  $\overline{H}_{(p)}^1(G) \neq 0$  for groups with infinitely many ends.

**Corollary 4.3.** *Let  $1 < p \in \mathbb{R}$ . If  $G$  is a finitely generated group with infinitely many ends, then  $\overline{H}_{(p)}^1(G) \neq 0$ .*

*Proof.* We will prove the corollary by constructing a nonconstant harmonic function, say  $h$ , on  $G$  that is an element of  $D^p(G)$ . It will then follow from the maximum principle for harmonic functions that  $h \notin C_0(G)$ . By the theorem  $h \notin (\overline{L^p(G)})_{D^p(G)}$  since  $L^p(G) \subset C_0(G)$ . Theorem 3.5 then shows  $h = u + v$ , where  $u \in \overline{L^p(G)}_{D^p(G)}$  and  $v$  is a nonconstant  $p$ -harmonic function, thus  $h$  will represent a nonzero class in  $\overline{H}_{(p)}^1(G)$ . We now proceed to construct  $h$  by using a technique that was used in the proof of Theorem 4.1 of [11].

Recall that  $S$  is a symmetric generating set for  $G$ . Define an element  $P$  in  $\mathbb{R}G$  by  $P = \frac{1}{|S|} \sum_{s \in S} s$ . Let  $f \in L^p(G)$  and define a bounded linear operator on  $L^p(G)$  by  $P(f) = f * P$ . Observe that  $f$  is a harmonic function on  $G$  if  $f * (P - I) = 0$ , where  $I$  is the identity operator on  $L^p(G)$ . Let  $k \in \mathbb{N}$  and denote by  $P^k$  multiplication of  $P$  with itself  $k$  times. Since  $G$  is nonamenable,  $\|P\| < 1$  in the operator norm for  $1 < p \in \mathbb{R}$  [4]. Thus  $F = -\sum_{k=0}^{\infty} P^k$  is a bounded operator on  $L^p(G)$ . Note  $F$  is the inverse of  $P - I$  in the space of bounded linear operators on  $L^p(G)$ . Let  $X$  denote the Cayley graph of  $G$  with respect to the generating set  $S$ . Thus the vertices of  $X$  are the elements of  $G$ , and  $g_1, g_2 \in G$  are joined by an edge if and only if  $g_1 = g_2 s^{\pm 1}$  for some generator  $s$ . Remove a finite number of vertices and edges of  $X$  to obtain two disjoint, infinite, connected components of  $X$ . Let  $X_1$  and  $X_2$  denote the components. Define a function  $h_1$  on  $G$  by

$$h_1(g) = \begin{cases} 2 & \text{if } g \text{ is a vertex in } X_1 \\ 1 & \text{if } g \text{ is a vertex in } X_2 \\ 0 & \text{otherwise} \end{cases}.$$

The support of  $h_1 * (P - I)$  is contained in  $\partial X_1 \cup \partial X_2$ , so  $h_1 * (P - I) \in \mathbb{R}G$ . Thus  $F(h_1 * (P - I)) \in L^p(G)$ . Set  $h_2 = F(h_1 * (P - I))$  and let  $h = h_1 - h_2$ . Note  $h \in D^p(G)$  since  $h_1 \in D^p(G)$  and  $h_2 \in L^p(G)$ . Now,  $h * (P - I) = h_1 * (P - I) - h_2 * (P - I) = 0$ , so  $h$  is a harmonic on  $G$ .  $\square$

Alain Valette has pointed out a simpler, but different, proof of the above corollary. We now proceed to give a quick sketch of his proof. By combining Lemma 2 with the remark after Proposition 1 of [1] we have that the dimension of  $H^1(G, \mathbb{R}G)$  over  $\mathbb{R}$  is the number of ends of  $G$  minus one. The corollary now follows from Theorem 4.1. Valette's proof also shows that the corollary is true for  $p = 1$ .

The situation becomes unclear if  $G$  is nonamenable with one end. For example,  $H^1(SL_n(\mathbb{Z}), L^2(SL_n(\mathbb{Z}))) = 0$  for  $n \geq 3$  since  $SL_n(\mathbb{Z})$  has property  $T$  when  $n \geq 3$  [5]. On the other hand, if  $G$  is a fundamental group of a closed Riemann surface of genus at least 2, then  $H^1(G, L^2(G)) \neq 0$  [2]. A good deal of information about  $H^1(G, L^2(G))$  can be found in [1].

## 5. A DESCRIPTION OF $H^1(G, L^p(G))$

In this section we will describe the nonzero elements of  $H^1(G, L^p(G))$  for groups that have polynomial growth of (precise) degree  $n > p$ . These results are a generalization of results from Section 6 of [10], where  $H^1(G, L^2(G))$  was discussed.

Let  $d > 1$ . We shall say that  $G$  satisfies condition  $S_d$  if there exists a constant  $C > 0$  such that  $\|f\|_{\frac{d}{d-1}} \leq C \|f\|_{D(1)}$  for all  $f \in \mathbb{R}G$ . If  $f \in \mathcal{F}(G)$  and  $t \geq 1$ , then  $f^t$  will denote the function  $f^t(g) = (f(g))^t$ . The following was proved in [10] but we include it here for completeness.

**Lemma 5.1.** *Let  $G$  be a finitely generated group and let  $t$  be a real number greater than or equal to 2. If  $f$  is a non-negative real-valued function in  $\mathcal{F}(G)$ , then*

$$\|f^t\|_{D(1)} \leq 2t \sum_{g \in G} f^{t-1}(g) \left( \sum_{s \in S} |(f * (s-1))(g)| \right).$$

*Proof.* Let  $g \in G$  and let  $s \in S$ . It follows from the Mean Value Theorem applied to  $x^t$  that  $(r^t - s^t) \leq t(r^{t-1} + s^{t-1})(r - s)$  where  $r$  and  $s$  are real numbers with  $0 \leq s \leq r$ . Thus  $|f^t(gs^{-1}) - f^t(g)| \leq t(f^{t-1}(g) + f^{t-1}(gs^{-1}))|f(gs^{-1}) - f(g)|$ . We now obtain  $\|f^t\|_{D(1)} = \sum_{g \in G} \sum_{s \in S} |f^t * (s-1)(g)| \leq t \sum_{g \in G} \sum_{s \in S} (f^{t-1}(g) + f^{t-1}(gs^{-1}))|f(gs^{-1}) - f(g)| = 2t \sum_{g \in G} \sum_{s \in S} f^{t-1}(g)|f(gs^{-1}) - f(g)|$   $\square$

The next proposition is a generalization of Proposition 6.2 of [10]. I would like to thank Thomas Schick for showing me this generalization.

**Proposition 5.2.** *Let  $2 \geq p \in \mathbb{R}$  and let  $d > p$ . If  $G$  satisfies condition  $S_d$ , then there is a constant  $C' > 0$  such that  $\|f\|_{\frac{pd}{d-p}} \leq C' \|f\|_{D(p)}$  for all  $f \in \mathbb{R}G$ .*

*Proof.* Set  $t = \frac{pd-p}{d-p}$ . By property  $S_d$ , Lemma 5.1 and Hölder's inequality we have (assuming without loss of generality that  $f$  is non-negative).

$$\begin{aligned} \|f^{\frac{pd-p}{d-p}}\|_{\frac{d}{d-1}} &\leq C \|f^{\frac{pd-p}{d-p}}\|_{D(1)} \\ &\leq 2C \left( \frac{pd-p}{d-p} \right) \sum_{g \in G} f^{\frac{d(p-1)}{d-p}}(g) \left( \sum_{s \in S} |(f * (s-1))(g)| \right) \\ &= 2C \left( \frac{pd-p}{d-p} \right) \sum_{g \in G} \sum_{s \in S} f^{\frac{d(p-1)}{d-p}}(g) |f(gs^{-1}) - f(g)| \\ &\leq 2C \left( \frac{pd-p}{d-p} \right) \|f^{\frac{d(p-1)}{d-p}}\|_{\frac{p}{p-1}} \|f\|_{D(p)}. \end{aligned}$$

Observe  $\|f^{\frac{pd-p}{d-p}}\|_{\frac{d}{d-1}} = \|f^{\frac{pd}{d-p}}\|_1^{\frac{d-1}{d}}$  and  $\|f^{\frac{d(p-1)}{d-p}}\|_{\frac{p}{p-1}} = \|f^{\frac{pd}{d-p}}\|_1^{\frac{p-1}{p}}$ . Substituting we obtain  $\|f^{\frac{pd}{d-p}}\|_1^{\frac{d-1}{d}} \leq C' \|f^{\frac{pd}{d-p}}\|_1^{\frac{p-1}{p}} \|f\|_{D(p)}$ . Dividing this inequality by  $\|f^{\frac{pd}{d-p}}\|_1^{\frac{p-1}{p}}$  and noting that  $\|f^{\frac{pd}{d-p}}\|_1^{\frac{d-p}{pd}} = (\|f\|_{\frac{pd}{d-p}})^{\frac{d-p}{pd}}$  will yield the claim in the proposition.  $\square$

We shall say that a group  $G$  satisfies condition  $(IS)_d$  if  $|A|^{d-1} < C|\partial A|^d$  for all finite subsets  $A$  of  $G$  and a positive constant  $C$ . Varopoulos proves the following proposition on page 224 of [12], also see Chapter 1.4 of [15].

**Proposition 5.3.** *A finitely generated group  $G$  satisfies condition  $(IS)_d$  for some  $d \geq 1$  if and only if it satisfies condition  $S_d$ .*

Our next result will show that each nonzero class in  $H^1(G, L^p(G))$ , where  $G$  is a group with polynomial growth of (precise) degree  $d > 2$ , can be represented by a function in  $L^{p'}(G)$  for some fixed real number  $p' > p$ .

**Theorem 5.4.** *Let  $G$  be a finitely generated group with polynomial growth of (precise) degree  $d$ . If  $d > p \geq 2$ , then each nonzero class in  $H^1(G, L^p(G))$  can be represented by a function from  $L^{\frac{pd}{d-p}}(G)$ .*

*Proof.* Varopoulos proves in the papers [13, 14] that  $G$  has polynomial growth of (precise) degree  $d$  if and only if  $G$  satisfies condition  $(IS)_d$ . Thus  $G$  also satisfies condition  $S_d$ .

Let  $1_G$  denote the constant function one on  $G$ . If  $1_G \in (\overline{\mathbb{R}G})_{D^p(G)}$ , then there exists a sequence  $f_n$  in  $\mathbb{R}G$  such that  $\|1_G - f_n\|_{D^p(G)} \rightarrow 0$  but  $\|f_n\|_{\frac{pd}{d-p}} \not\rightarrow 0$  contradicting Proposition 5.2. Hence  $(\overline{\mathbb{R}G})_{D^p(G)} \neq D^p(G)$ . By Theorem 3.5 each  $f \in D^p(G)$  can be represented uniquely by  $u + h$ , where  $u \in (\overline{\mathbb{R}G})_{D^p(G)}$  and  $h \in HD^p(G)$ . By [8, Corollary 1.10],  $HD^p(G) = \mathbb{R}$ . Thus nonzero classes in  $H^1(G, L^p(G))$  can be represented by functions in  $(\overline{\mathbb{R}G})_{D^p(G)} \setminus L^p(G)$ . Let  $f \in (\overline{\mathbb{R}G})_{D^p(G)}$ , so there exists a sequence  $\{f_n\}$  in  $\mathbb{R}G$  such that  $f_n \rightarrow f$  in the Banach space  $D^p(G)$ . Observe that  $f_n \rightarrow f$  in  $D^r(G)$  for  $p \leq r < \infty$ . By Proposition 5.2  $\{f_n\}$  forms a Cauchy sequence in  $L^{\frac{pd}{d-p}}(G)$ . Let  $\bar{f}$  be the limit of this sequence in  $L^{\frac{pd}{d-p}}(G)$ . It now follows  $f_n \rightarrow \bar{f}$  in  $D^{\frac{pd}{d-p}}(G)$  since  $\|(\bar{f} - f_n) * (s - 1)\|_{\frac{pd}{d-p}} \rightarrow 0$  as  $n \rightarrow \infty$  for each  $s \in S$ . Therefore  $\bar{f} = f$  since  $f_n \rightarrow f$  in  $D^{\frac{pd}{d-p}}(G)$ .  $\square$

**Corollary 5.5.** *Let  $2 < p \in \mathbb{R}$  and let  $d$  be an integer greater than  $p$ . Each nonzero class in  $H^1(\mathbb{Z}^d, L^p(\mathbb{Z}^d))$  can be represented by a function from  $L^{\frac{pd}{d-p}}(\mathbb{Z}^d)$ .*

## 6. SOME INCLUSION RESULTS

Since  $L^p(G) \subseteq L^{p'}(G)$  for  $1 < p \leq p' \in \mathbb{R}$ , a natural question to ask is how does  $H^1(G, L^p(G))$  relate to  $H^1(G, L^{p'}(G))$ . In this section we will give some answers to this question.

**Lemma 6.1.** *Let  $h \in HD^p(G)$ . If  $h \in C_0(G)$ , then  $h = 0$ .*

*Proof.* The set  $\{g \mid |h(g)| > \epsilon\}$  is finite for a given  $\epsilon > 0$ . Thus there exists an  $x \in G$  such that  $|h(x)| \geq |h(g)|$  for all  $g \in G$ . It follows  $h(x) = h(xs^{-1})$  for all  $s \in S$  since  $\sum_{s \in S} |h(xs^{-1}) - h(x)|^{p-2} h(xs^{-1}) = \sum_{g \in S} |h(xs^{-1}) - h(x)|^{p-2} h(x)$ . We now obtain  $h(x) = h(g)$  for all  $g \in G$  since the Cayley graph of  $G$  is connected. Therefore  $h = 0$ .  $\square$

We now give an inclusion result for nonamenable groups.

**Proposition 6.2.** *Let  $G$  be a finitely generated nonamenable group. If  $1 < p \leq p' \in \mathbb{R}$ , then  $H^1(G, L^p(G)) \subseteq H^1(G, L^{p'}(G))$ .*

*Proof.* Let  $f$  represent a nonzero class in  $H^1(G, L^p(G))$ . Thus  $f \in D^p(G) \subseteq D^{p'}(G)$  and  $f \notin L^p(G)$ . By Theorem 3.5 we can uniquely write  $f = u + h$ , where  $u \in L^p(G)$  and  $h$  is a nonconstant element in  $HD^p(G)$ . By Lemma 6.1  $h \notin C_0(G)$  so it follows  $f \notin L^{p'}(G)$ . Hence  $f$  also represents a nonzero class in  $H^1(G, L^{p'}(G))$ .  $\square$

We now finish this section by giving an example to show the above proposition is not true for amenable groups. Define  $f: \mathbb{Z} \rightarrow \mathbb{R}$  by

$$f(n) = \begin{cases} \frac{1}{\sqrt[4]{n}} & n \geq 1 \\ 0 & \text{otherwise} \end{cases}$$



for some  $1 < p \in \mathbb{R}$ . Observe that  $f \notin L^p(\mathbb{Z})$  but  $f \in L^{p'}(\mathbb{Z})$  for  $p' > p$ . Now

$$\begin{aligned} \sum_{n=1}^{\infty} |(f * (s-1))(n)|^p &= \sum_{n=1}^{\infty} |f(n-1) - f(n)|^p \\ &= 1 + \sum_{n=2}^{\infty} \left| \frac{\sqrt[p]{n} - \sqrt[p]{n-1}}{\sqrt[p]{n(n-1)}} \right|^p \\ &\leq 1 + \sum_{n=2}^{\infty} \frac{1}{(n-1)^2}. \end{aligned}$$

Thus  $f \in D^p(\mathbb{Z})$  which implies  $f$  represents a nonzero class in  $H^1(\mathbb{Z}, L^p(\mathbb{Z}))$  but  $f$  is in the zero class of  $H^1(\mathbb{Z}, L^{p'}(\mathbb{Z}))$  for  $p' > p$ .

#### REFERENCES

- [1] Mohammed E. B. Bekka and Alain Valette. Group cohomology, harmonic functions and the first  $L^2$ -Betti number. *Potential Anal.*, 6(4):313–326, 1997.
- [2] Jeff Cheeger and Mikhael Gromov.  $L_2$ -cohomology and group cohomology. *Topology*, 25(2):189–215, 1986.
- [3] Ivar Ekeland and Roger Témam. *Convex analysis and variational problems*, volume 28 of *Classics in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1999.
- [4] Peter Gerl. Random walks on graphs with a strong isoperimetric property. *J. Theoret. Probab.*, 1(2):171–187, 1988.
- [5] A. Guichardet. Étude de la  $l$ -cohomologie et de la topologie du dual pour les groupes de Lie à radical abélien. *Math. Ann.*, 228(3):215–232, 1977.
- [6] Ilkka Holopainen. Rough isometries and  $p$ -harmonic functions with finite Dirichlet integral. *Rev. Mat. Iberoamericana*, 10(1):143–176, 1994.
- [7] Ilkka Holopainen and Paolo M. Soardi.  $p$ -harmonic functions on graphs and manifolds. *Manuscripta Math.*, 94(1):95–110, 1997.
- [8] Ilkka Holopainen and Paolo M. Soardi. A strong Liouville theorem for  $p$ -harmonic functions on graphs. *Ann. Acad. Sci. Fenn. Math.*, 22(1):205–226, 1997.
- [9] C. H. Houghton. Ends of groups and the associated first cohomology groups. *J. London Math. Soc. (2)*, 6:81–92, 1972.
- [10] Michael J. Puls. Group cohomology and  $L^p$ -cohomology of finitely generated groups. *Canad. Math. Bull.*, 46(2):268–276, 2003.
- [11] Paolo M. Soardi and Wolfgang Woess. Uniqueness of currents in infinite resistive networks. *Discrete Appl. Math.*, 31(1):37–49, 1991.
- [12] N. Th. Varopoulos. Isoperimetric inequalities and Markov chains. *J. Funct. Anal.*, 63(2):215–239, 1985.
- [13] Nicholas Th. Varopoulos. Théorie du potentiel sur les groupes nilpotents. *C. R. Acad. Sci. Paris Sér. I Math.*, 301(5):143–144, 1985.
- [14] Nicholas Th. Varopoulos. Théorie du potentiel sur des groupes et des variétés. *C. R. Acad. Sci. Paris Sér. I Math.*, 302(6):203–205, 1986.
- [15] Wolfgang Woess. *Random walks on infinite graphs and groups*, volume 138 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2000.
- [16] Maretsugu Yamasaki. Ideal boundary limit of discrete Dirichlet functions. *Hiroshima Math. J.*, 16(2):353–360, 1986.

DEPARTMENT OF MATHEMATICS, EASTERN OREGON UNIVERSITY, ONE UNIVERSITY BOULEVARD, LA GRANDE, OR 97850, USA

*E-mail address:* mpuls@eou.edu